

CLIENT INFORMATION LEAFLET

Updated: July 2019

Trazodone

- **Keep this leaflet safe, as you may need to refer to it again.**
- **Please ask your vet or veterinary nurse if you have any further questions.**
- **This medicine has been prescribed for *your* pet ONLY. Do not take it yourself or give it to another person or any other animal; it may harm them even if their symptoms appear to be the same.**

The medicine you have been given for your dog or cat is called trazodone. It may have a trade name such as Desyrel®, Molipaxin® or Oleptro®, but often will just be called trazodone.

What is trazodone?

Trazodone is an antidepressant medication within a class of medications called serotonin modulators. It works by controlling the levels of a natural substance in the brain called serotonin, which is involved in the regulation of mood, anxiety, sleep, appetite and energy levels.

Why has my pet been prescribed trazodone?

In dogs, trazodone is used for chronic anxiety-related disorders that are unresponsive to other medications. The anti-anxiety and mood regulating properties of this drug make it particularly suitable for the treatment of separation anxiety in association with a behaviour modification plan. In cats, trazodone may be used for the management of short-term anxiety-related problems such as travel anxiety.

How should I store trazodone?

This medicine does not require any special storage conditions. Store in a cool, dry place. For safety, **all medicines should be kept out of the reach and sight of children.**

How do I give trazodone tablets to my pet?

This medicine may be given either with or without food. The tablets may be disguised in a small amount of strongly flavoured food that your pet likes. Alternatively, they can be placed carefully on the back of your pet's tongue and their mouth closed until the entire tablet has been swallowed. Treatment should be given as directed by your vet but is often given as an oral treatment once a day.

How long will my pet need to take trazodone?

The duration of administration depends on the condition being treated, the response to medication and the development of any adverse side effects. In certain situations, the dose may be increased after a few weeks until an appropriate unique dose is found

for your pet. Your vet will advise you on the length of time for which you will need to give this medicine. This may vary between patients and can be for the rest of their life, although increased tolerance to this medication over time is common.

What should I do if I run out of tablets?

Try not to run out. Make sure you order more tablets from your vet if your supply is getting low. If you do run out, contact your own vet for further advice and restart the course as soon as possible.

What should I do if I miss a dose?

If a dose is missed give the medication as soon as possible and continue the regular schedule. **DO NOT** give a double dose to make up for the missed dose and do not exceed the total stated dose in any one 24-hour period.

What should I do if my pet is accidentally given too many doses?

Contact your vet immediately if an overdose is given. The most common side effects with an overdose are lethargy, drooling, agitation and in some cases, seizures.

Can my pet take trazodone if I am already giving them other drugs?

Tell your vet if you are giving your pet any other medications, even if you think they already know. This includes herbal and off-the-shelf remedies from a pet shop, pharmacy or the internet. This drug interacts with a wide range of medications and your vet will advise you on the safety of taking multiple medications accordingly. In particular, trazodone should not be used alongside some other antidepressant medications, especially in the class of 'monoamine oxidase inhibitors' (e.g. selegine). In some cases, these medications should not be given within a few weeks of one another. Although trazodone may be given alongside certain other anti-anxiety medications in low doses, caution should be taken as an additive

effect is possible. Particular caution should be taken if trazodone is given with antacid medications such as cimetidine, certain anti-seizure medications such as phenobarbital as well as diuretics. If taken with blood thinners such as warfarin and aspirin, prolonged bleeding may occur. Certain medications used to treat fungal diseases such as ketoconazole and itraconazole can prolong the effects of trazodone. If you have any concerns about using trazodone with any other medications, please ask your vet for further advice.

What are the possible side effects of trazodone for my pet?

Side effects include sedation, drowsiness, vomiting, excitability and a dry mouth. Weakness, confusion, restlessness and irritability may also be caused. In humans, persistent painful erections have been reported while taking trazodone. If your pet experiences an allergic reaction to the medication, side effects may include facial swelling, hives, vomiting and seizures.

What should I do if my pet is unwell while taking trazodone?

If your pet is unwell while receiving this medication, you should not give any further doses and should contact your vet immediately for advice.

What should I do if a person accidentally takes this drug?

If a person takes your pet's tablets, the person should be taken to the local hospital immediately. Take this leaflet and any remaining tablets plus their container (even if it is empty) with you.

Whom do I contact if I want to know more?

If you have any questions about this drug, or any concerns with your pet's health then contact your own vet. They will know your pet's medical history and will know about trazodone.

The Prescribing Cascade

This medicine is authorized for use in human patients and is used by vets under the 'prescribing cascade'. The medicine is not authorized by the Veterinary Medicines Directorate (VMD), an executive agency of the Department for Environment, Food and Rural Affairs (Defra), for use in dogs/cats/pets. Your vet can explain the 'prescribing cascade' in further detail to you and also explain why they are prescribing this drug for your pet. You will be asked to sign a consent form stating that you understand the reasons that the drug is being prescribed and its possible complications, before the treatment is issued.

© British Small Animal Veterinary Association 2019. While the editors and the BSAVA have made every effort in preparing this information leaflet, the contents and any statements are made in good faith purely for general guidance and cannot be regarded as substitute for professional advice. The publishers, contributors and the BSAVA do not take responsibility for the information provided on this leaflet and hence do not accept any liability for loss or expense incurred (by you or persons that you disseminate the materials to) as a result of relying on content in this leaflet. To this end, you are advised to consult your vet and seek their professional advice before taking any steps set out in this leaflet. If you are a vet, you must not rely on the contents in this leaflet without independently verifying the correctness and veracity of the contents. BSAVA is not responsible for any alterations made to this document from the version supplied.